

## NC IMMUNIZATION REQUIREMENTS 2019-2020

<b>Pre-K</b>	<b>Grades K-4</b>	<b>Grades 5-6</b>	<b>Grades 7-11</b>	<b>Grade 12</b>
<b>(4) DTP/DTaP</b>	<b>*(5) DTP/DTaP</b> with booster on/after 4 <sup>th</sup> birthday	<b>*(5) DTP/DTaP</b> with booster on/after 4 <sup>th</sup> birthday	<b>*(5) DTP/DTaP</b> with booster on/after 4 <sup>th</sup> birthday	<b>*(5) DTP/DTaP</b> with booster on/after 4 <sup>th</sup> birthday
			<b>Tdap</b> Required if individual had not previously received it and are entering the 7 <sup>th</sup> grade or by 12 years of age whichever comes first. Effective 7-1-2015	<b>Tdap</b> (Booster required if entering 6 <sup>th</sup> grade on/after 8-1-2008, if 5 years or more have passed since the last dose of tetanus/diphtheria toxoid)
<b>(3) Polio</b>	<b>** (4) Polio</b> 4 <sup>th</sup> dose is required on/after the 4 <sup>th</sup> birthday and before entering school for the first time. Effective 7-1-2015	<b>*** (4) Polio</b>	<b>*** (4) Polio</b>	<b>*** (4) Polio</b>
<b>**** (1) MMR</b> On/after 1 <sup>st</sup> birthday	<b>**** (2) MMR</b> (1 <sup>st</sup> dose on/after 1 <sup>st</sup> birthday) At least 28 day interval needed between doses	<b>**** (2) MMR</b> (1 <sup>st</sup> dose on/after 1 <sup>st</sup> birthday) At least 28 day interval needed between doses	<b>**** (2) MMR</b> (1 <sup>st</sup> dose on/after 1 <sup>st</sup> birthday) At least 28 day interval needed between doses	<b>**** (2) MMR</b> (1 <sup>st</sup> dose on/after 1 <sup>st</sup> birthday) At least 28 day interval needed between doses
<b>(3) Hepatitis-B</b> Third dose shall NOT be given prior to 24 weeks of age if born on/after 11-17-2001	<b>(3) Hepatitis-B</b> Third dose shall NOT be given prior to 24 weeks of age if born on/after 11-17-2001	<b>(3) Hepatitis-B</b> Third dose shall NOT be given prior to 24 weeks of age if born on/after 11-17-2001	<b>(3) Hepatitis-B</b> Third dose shall NOT be given prior to 24 weeks of age if born on/after 11-17-2001	<b>(3) Hepatitis-B</b> Third dose shall NOT be given prior to 24 weeks of age if born on/after 11-17-2001
<b>(1) Varicella</b> On/after 1 <sup>st</sup> birthday	<b>(2) Varicella</b> (1 <sup>st</sup> dose on/after 1 <sup>st</sup> birthday) At least 28 day interval needed between doses. (As of 7-1-2015, historical documentation of disease must include approximate date or age of infection and a health care provider signature.)	<b>(1) Varicella</b> (1 <sup>st</sup> dose on/after 1 <sup>st</sup> birthday) Required if born on/after 4-1-2001 and no varicella disease documentation. Parent or health care provider may provide written history of disease but must include approximate date or age of infection.	<b>(1) Varicella</b> (1 <sup>st</sup> dose on/after 1 <sup>st</sup> birthday) Required if born on/after 4-1-2001 and no varicella disease documentation. Parent or health care provider may provide written history of disease but must include approximate date or age of infection.	<b>(1) Varicella</b> (1 <sup>st</sup> dose on/after 1 <sup>st</sup> birthday) Required if born on/after 4-1-2001 and no varicella disease documentation. Parent or health care provider may provide written history of disease but must include approximate date or age of infection.
<b>(1-4) Hib</b> Hib vaccine is not given after 5 years of age. If the first dose is given on/after 15 months of age, series is complete. If 4 <sup>th</sup> dose is given on/after 12 months of age, series is complete. (Dosage requirement may vary depending on the vaccine type and the age when received first dose.)	<b>Hib</b> Hib vaccine is not given after 5 years of age. If not yet 5 years of age, same requirements as for Pre-K.			
<b>(1-4) Pneumococcal Conjugate</b> Is required for individuals born on/after 7-1-2015 and if younger than 5 years of age. If the first dose is given on/after 24 months of age, series is complete. (Dosage and interval requirements may vary depending on the age when received first dose.)			<b>(1) Meningococcal conjugate</b> (1 <sup>st</sup> dose on/after 10 <sup>th</sup> birthday) Required for individuals born on/after 1-1-2003 and entering the 7 <sup>th</sup> grade or by 12 years of age, whichever comes first. (Effective 7-1-2015)	

NOTE: Four-day rule applies to all required minimum ages.

\* Children receiving #4 DTP/DTaP on or after 4<sup>th</sup> birthday are not required to have #5.

\*\* Children receiving #3 Polio on or after 4<sup>th</sup> birthday and at least 6 months after the 2<sup>nd</sup> dose are not required to have #4.

\*\*\* Children receiving #3 Polio on or after their 4<sup>th</sup> birthday are not required to have #4. Children 18 years or older are not required to receive a polio vaccine.

\*\*\*\* If given separately: Two measles doses, two mumps doses, and one rubella dose required. First dose must be on or after first birthday and at least a 28 day interval is needed between the first and second doses. Second mumps dose only required if entered school on or after 7-1-2008.